

Physicians Laboratory Services Cumulative Antimicrobial Susceptibility Report
All Clients, Other Isolates | January 2021 - December 2021

% Susceptible Gram Positive	Total*	Penicillin(2)	Ampicillin	Oxacillin(1)	Amoxicillin/Clavulanate	Cefotaxime(2)	Ceftriaxone(2)	Meropenem	Gentamicin	Ciprofloxacin(7)	Levofloxacin(7)	Moxifloxacin	Trimethoprim/Sulfa	Clindamycin(5)	Daptomycin(6)	Erythromycin(5)	Nitrofurantoin(3)	Linezolid(4)	Vancomycin	Tetracycline(7)	Gentamicin Synergy(8)	Streptomycin Synergy(8)	Penicillin (non-CSF)(2)	Penicillin V (oral)(2)	Cefotaxime (non-CSF)(2)	Ceftriaxone (non-CSF)(2)
<i>Staph. aureus</i> (MSSA) 71%	560	-	-	100	-	-	-	-	99	87	89	96	99	79	100	66	-	100	100	92	-	-	-	-	-	-
<i>Staph. aureus</i> (MRSA) 29%	226	--	--	0	--	--	--	--	98	43	46	77	96	72	98	18	--	100	100	89	--	--	--	--	--	--
<i>Staph.</i> spp. (coagulase-negative)	118	--	--	69	--	--	--	--	97	81	83	94	81	71	100	54	--	100	100	87	--	--	--	--	--	--
<i>Strep. pneumoniae</i>	18	65	--	--	82	82	88	76	--	--	100	--	71	79	--	53	--	--	100	71	--	--	100	58	100	100
<i>Enterococcus faecalis</i>	169	100	100	--	--	R	R	--	--	82	86	--	R	R	100	--	--	99	99	31	--	--	--	--	--	--
<i>Enterococcus faecium</i>	6	33	33	--	--	R	R	--	--	17	33	--	R	R	100	--	--	100	33	33	--	--	--	--	--	--
<i>Enterococcus</i> spp.	13	92	92	--	--	R	R	--	--	69	69	--	R	R	100	--	--	100	54	62	--	--	--	--	--	--

The % susceptible for each organism/antimicrobial combination was generated by including the first isolate of that organism encountered on a given patient.

Other Isolates includes isolates from body sites other than urine and blood such as wound, tissue, genital, body fluid, and respiratory.

R Organism has intrinsic resistance to this antimicrobial.

- Not tested/indicated for organism.

* Antibiograms created for organisms with less than 30 total isolates have questionable statistical significance. Interpret data with caution.

(1) Oxacillin predicts susceptibility to most cephalosporins, carbapenems, and beta-lactam/beta-lactamase inhibitors.

(2) For *S. pneumoniae*: cefotaxime, ceftriaxone, and penicillin % susceptible listed is based on parenteral CSF (meningitis), parenteral non-CSF, and oral MIC breakpoints.

(3) Nitrofurantoin is reported for isolates from urine only.

(4) Linezolid is reported for isolates from non-urine sources only.

(5) For *Staphylococcus* spp.: clindamycin and erythromycin are reported for isolates from non-urine sources only.

(6) For *Staphylococcus* spp.: daptomycin is reported for isolates from non-respiratory sources only.

(7) For all *Enterococcus* spp.: ciprofloxacin, levofloxacin, and tetracycline are reported for isolates from urine only.

(8) For *E. faecalis* and *E. faecium*: gentamicin and streptomycin high-level resistance testing is reported for isolates from blood cultures only.

