

Physicians Laboratory Services Cumulative Antimicrobial Susceptibility Report
All Clients, Blood/Systemic Isolates | January 2022 - December 2022

% Susceptible Gram Positive	Total*	Penicillin(2)	Ampicillin	Oxacillin(1)	Amoxicillin/Clavulanate	Cefotaxime(2)	Ceftriaxone(2)	Meropenem	Gentamicin	Ciprofloxacin(7)	Levofloxacin(7)	Moxifloxacin	Trimethoprim/Sulfa	Clindamycin(5)	Daptomycin(6)	Erythromycin(5)	Nitrofurantoin(3)	Linezolid(4)	Vancomycin	Tetracycline(7)	Gentamicin Synergy(8)	Streptomycin Synergy(8)	Penicillin (non-CSF)(2)	Cefotaxime (non-CSF)(2)	Ceftriaxone (non-CSF)(2)	
<i>Staph. aureus</i> (MSSA) 79%	15	--	--	100	--	--	--	--	100	87	87	100	100	67	100	53	--	100	100	93	--	--	--	--	--	
<i>Staph. aureus</i> (MRSA) 21%	4	--	--	0	--	--	--	--	100	0	0	50	100	75	100	0	--	100	100	75	--	--	--	--	--	
<i>Staph.</i> spp. (coagulase-negative)	8	--	--	63	--	--	--	--	50	25	25	50	25	50	100	38	--	100	100	88	--	--	--	--	--	
<i>Strep. pneumoniae</i>	7	86	--	--	100	100	100	86	--	--	100	--	100	100	--	71	--	--	100	100	--	--	100	80	100	100
<i>Enterococcus faecalis</i>	7	100	100	--	--	R	R	--	--	86	86	--	R	R	100	--	--	100	100	29	86	86	--	--	--	
<i>Enterococcus faecium</i>	0	--	--	--	--	R	R	--	--	--	--	--	R	R	--	--	--	--	--	--	--	--	--	--	--	
<i>Enterococcus</i> spp.	1	100	100	--	--	R	R	--	--	100	100	--	R	R	100	--	--	100	0	100	--	--	--	--	--	

The % susceptible for each organism/antimicrobial combination was generated by including the first isolate of that organism encountered on a given patient.

R Organism has intrinsic resistance to this antimicrobial.

- Not tested/indicated for organism.

* Antibiograms created for organisms with less than 30 total isolates have questionable statistical significance. Interpret data with caution.

(1) Oxacillin predicts susceptibility to most cephalosporins, carbapenems, and beta-lactam/beta-lactamase inhibitors.

(2) For *S. pneumoniae*: cefotaxime, ceftriaxone, and penicillin % susceptible listed is based on parenteral CSF (meningitis), parenteral non-CSF, and oral MIC breakpoints.

(3) Nitrofurantoin is reported for isolates from urine only.

(4) Linezolid is reported for isolates from non-urine sources only.

(5) For *Staphylococcus* spp.: clindamycin and erythromycin are reported for isolates from non-urine sources only.

(6) For *Staphylococcus* spp.: daptomycin is reported for isolates from non-respiratory sources only.

(7) For all *Enterococcus* spp.: ciprofloxacin, levofloxacin, and tetracycline are reported for isolates from urine only.

(8) For *E. faecalis* and *E. faecium*: gentamicin and streptomycin high-level resistance testing is reported for isolates from blood cultures only.

Physicians Laboratory Services Cumulative Antimicrobial Susceptibility Report
All Clients, Blood/Systemic Isolates | January 2022 - December 2022

% Susceptible Gram Negative	Total*	Ampicilin	Ampicillin/Sulbactam	Piperacillin/Tazobactam	Cefazolin	Ceftazidime	Ceftazidime/Avibactam	Ceftriaxone	Cefepime	Cefoxitin	Aztreonam	Ertapenem	Imipenem	Meropenem	Amikacin	Gentamicin	Tobramycin	Ciprofloxacin(3)	Levofloxacin(3)	Trimethoprim/Sulffa	Nitrofurantoin(1)	Tetracycline(2)	Tigecycline
<i>Escherichia coli</i>	16	69	69	100	94	94	100	94	94	100	94	100	100	100	100	100	100	81	81	88	--	94	100
<i>Klebsiella pneumoniae</i>	7	R	86	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	--	86	100
<i>Proteus mirabilis</i>	1	100	100	100	100	100	100	100	100	100	100	100	100	--	100	100	100	100	100	100	R	R	--
<i>Klebsiella (Enterobacter) aerogenes</i>	3	R	R	100	R	33	100	33	100	R	33	100	67	100	100	100	100	100	100	100	--	67	100
<i>Enterobacter cloacae</i> complex	3	R	R	100	R	100	100	67	100	R	100	100	100	100	100	100	100	100	100	100	--	100	100
<i>Serratia marcescens</i>	1	R	R	100	R	100	100	100	100	R	100	100	--	100	100	100	100	100	100	100	R	0	100
<i>Pseudomonas aeruginosa</i>	4	R	R	100	R	100	100	R	100	R	100	R	75	100	100	100	100	75	100	R	--	R	--
<i>Acinetobacter baumannii</i>	0	R	--	--	R	--	--	--	--	R	R	R	--	--	--	--	--	--	--	--	--	--	
<i>Citrobacter freundii</i> complex	1	R	R	100	R	100	100	100	100	R	100	100	100	100	100	100	100	100	100	100	--	100	100
<i>Morganella morganii</i>	0	R	--	--	R	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	R	R	--
<i>Providencia</i> spp.	0	R	--	--	R	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	R	R	--
<i>Salmonella</i> spp.	3	100	--	--	--	100	--	--	--	--	--	--	--	--	--	--	--	100	100	100	--	--	--
<i>Shigella</i> spp.	0	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
<i>Stenotrophomonas</i>	0	R	R	R	R	--	--	R	--	R	R	R	R	R	R	R	R	--	--	--	--	R	--
<i>Klebsiella oxytoca</i>	1	R	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	--	100	100
<i>Haemophilus influenzae</i>	0	Beta-lactamase positive: --%																					

The % susceptible for each organism/antimicrobial combination was generated by including the first isolate of that organism encountered on a given patient.

R Organism has intrinsic resistance to this antimicrobial.

-- Not tested/indicated for organism.

* Antibiograms created for organisms with less than 30 total isolates have questionable statistical significance. Interpret data with caution.

(1) Nitrofurantoin is reported for isolates from urine only.

(2) For *A. baumannii*: tetracycline is reported for isolates from urine only.

(3) Ciprofloxacin/Levofloxacin: Enterobacteriales & *P. aeruginosa* %Susceptible utilizes CLSI M100 28th Edition breakpoints.